

AD-A071 321

SRI INTERNATIONAL MENLO PARK CA

F/G 4/1

SCINTILLATION STRUCTURE IN A SEVERELY DISTURBED SCATTERING ENVI--ETC(U)

OCT 78 C L RINO

DNA001-77-C-0038

UNCLASSIFIED

DNA-4732T

NL

| OF |
AD
A071321



12

LEVEL III

AD-E300532

DNA 4732T

DA071321

SCINTILLATION STRUCTURE IN A SEVERELY DISTURBED SCATTERING ENVIRONMENT

Charles L. Rino
SRI International
333 Ravenswood Avenue
Menlo Park, California 94025

October 1978

Topical Report 2 for Period 1 November 1977-30 September 1978

CONTRACT No. DNA 001-77-C-0038

APPROVED FOR PUBLIC RELEASE;
DISTRIBUTION UNLIMITED.

THIS WORK SPONSORED BY THE DEFENSE NUCLEAR AGENCY
UNDER RDT&E RMSS CODE B322078464 S99QAXHB05415 H2590D.

Prepared for
Director
DEFENSE NUCLEAR AGENCY
Washington, D. C. 20305

DDC
RECEIVED
JUL 18 1979
B

DDC FILE COPY

79 05 18 016

Destroy this report when it is no longer
needed. Do not return to sender.

PLEASE NOTIFY THE DEFENSE NUCLEAR AGENCY,
ATTN: TISI, WASHINGTON, D.C. 20305, IF
YOUR ADDRESS IS INCORRECT, IF YOU WISH TO
BE DELETED FROM THE DISTRIBUTION LIST, OR
IF THE ADDRESSEE IS NO LONGER EMPLOYED BY
YOUR ORGANIZATION.



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER DNA 4732T	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) SCINTILLATION STRUCTURE IN A SEVERELY DISTURBED SCATTERING ENVIRONMENT		5. TYPE OF REPORT & PERIOD COVERED Topical Report 2 for Period 1 Nov 77-30 Sep 78
7. AUTHOR(s) Charles L. Rino		6. PERFORMING ORG. REPORT NUMBER SRI Project 5960
9. PERFORMING ORGANIZATION NAME AND ADDRESS SRI International 333 Ravenswood Avenue Menlo Park, California 94025		8. CONTRACT OR GRANT NUMBER(s) DNA 001-77-C-0038
11. CONTROLLING OFFICE NAME AND ADDRESS Director Defense Nuclear Agency Washington, D.C. 20305		10. PROGRAM ELEMENT PROJECT, TASK AREA & WORK UNIT NUMBERS Subtask S99QAXHB054-15
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE October 1978
		13. NUMBER OF PAGES 38
		15. SECURITY CLASS (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES This work sponsored by the Defense Nuclear Agency under RDT&E RMSS Code B322078464 S99QAXHB05415 H2590D.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Scintillation Striations Phase-Screen Model Strong-Scatter		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This report presents a complete treatment of the second-order statistics of intensity under conditions of strong scattering. A three-dimensional powerlaw striation model is used in the analysis. For completeness, the special forms for a strictly two-dimensional medium are also derived. Asymptotic formulas for the fade coherence time are derived. Simple asymptotic results can be derived only for three-dimensional spectral indices less than 4. However, naturally occurring striations seem to fall in this range, and the simple formulas are generally applicable.		

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED.

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

CONTENTS

LIST OF ILLUSTRATIONS	2
LIST OF TABLES	2
I EXECUTIVE SUMMARY	3
II INTRODUCTION	5
III GENERAL RESULTS FOR A GAUSSIAN PHASE SCREEN	8
IV SCATTER IN STRICT POWER-LAW ENVIRONMENTS	12
V ASYMPTOTIC RESULTS FOR LARGE AND SMALL VALUES OF U	22
VI SUMMARY AND DISCUSSION	26
REFERENCES	31

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or special
A	

ILLUSTRATIONS

1	Ratio of S_4 to U vs ν Showing Discontinuity at $\nu = 1.5$	21
2	Curves of Constant S_4 on a Plot of $\lambda z/r_o^2$ vs φ_o --Taken from Singleton (1970)	30

TABLE

1	Limiting Forms of $g(\vec{\xi}, \vec{\eta})$ and $g'(\zeta, \eta)$	16
---	--	----

I EXECUTIVE SUMMARY

In severely disturbed propagation environments, the scintillation structure asymptotically approaches a form that can be characterized by simple formulas. For example, the intensity statistics are accurately approximated by a Rayleigh distribution. Since the performance of both coherent and non-coherent systems in a Rayleigh fading environment is critically dependent on the fade coherence time, the second-order statistics must also be specified to complete the phenomenology-systems analysis link.

This report presents a complete treatment of the second-order statistics of intensity in a power-law scattering medium. A three-dimensional striation model is used, so that the propagation angle dependence in a three-dimensional anisotropic medium can be properly evaluated. For completeness, however, the special forms that apply to a strictly two-dimensional medium are also included. Thus, the results of the study are applicable to the Wideband satellite data base as well as the various numerical simulations that have been performed.

An important finding from this study is that the asymptotic behavior of the second-order statistics of intensity under strong scatter conditions is critically dependent on the power-law index. The theory admits three-dimensional striation spectral densities of the form $q^{-\alpha}$ where $2 < \alpha < 6$. When $\alpha < 4$, the contribution of large-scale structures is strongly suppressed by Fresnel filtering. When $\alpha \geq 4$, the large-scale structures are only weakly suppressed, and that the strong-scatter behavior begins to show some characteristics of scattering in a medium that contains a single dominant scale size.

When $\alpha < 4$, the second-order statistics asymptotically approach a form that is independent of the Fresnel radius, and a simple expression for the fade coherence time can be obtained. In light of data from the

Wideband satellite, it appears that the effective three-dimensional spectral index is indeed somewhat less than 4. Thus, the simpler asymptotic results are applicable. When $\alpha \geq 4$, the intensity statistics retain a dependence on the Fresnel radius, and simple limiting forms cannot be obtained.

A detailed summary of the results is given in Section VI as is a comparison with earlier results based on a gaussian spectral density function. Under conditions of weak scatter, the results converge to the weak-scatter forms that were independently derived in Rino and Matthews (1978). The intermediate range between weak and strong scattering must be treated by numerical computations or more elaborate asymptotic approximations.

II INTRODUCTION

A full treatment of multiple scattering in an extended three-dimensional medium is now well formulated [see, for example, Rino (1978) and the references cited therein]. However, it is impractical to solve the necessary vector differential equations numerically except in certain special cases. Thus, simplified models and/or asymptotic results are invariably used to interpret data. Fante (1975) has reviewed the various asymptotic results that have been applied to laser beam propagation where isotropic scattering is an appropriate idealization.

Ionospheric scattering, by comparison, is complicated by the fact that the irregularities are highly anisotropic. Moreover, the ionospheric outer-scale cutoff, q_0 , is sufficiently small that the condition $q_0/Z \ll 1$, where $Z = \lambda \tilde{Z} \sec \theta / (4\pi)$ is the Fresnel area, holds for all observing geometries at the lowest frequencies of interest ($\lambda \leq 3m$). This fact can be used to greatly simplify the theory.

In Rino and Matthews (1978) (hereafter Paper I), the weak-scatter theory was reformulated to show explicitly the ramifications of the very large outer scale $\ell_0 = 2\pi/q_0$. In particular, a closed form analytic expression for the intensity scintillation index S_4 was derived by taking the limit of the integral expression as q_0 approaches zero. The resulting expression is valid when $q_0/Z \ll 1$, which is a very good approximation as noted above.

In this paper, the results for the second-order statistics of intensity are extended to accommodate strong scattering. The analysis is based on the formulation of the gaussian phase-screen model developed by Gochelashvily and Shishov (1971). The general results are reviewed in Section III, where it is shown that, to calculate the second-order statistics of intensity, only the phase-structure function need be specified.

In Section IV of this paper, the limiting forms of the general results are computed as q_0 approaches zero, as was done in Paper I for weak scatter. The analysis is identical to that presented by Rumsey (1975). For completeness, however, the special forms for a one-dimensional phase screen are also included. The scattering from a power-law phase screen is governed by a single "universal" strength parameter, U , which combines the effects of perturbation strength and distance from the scattering medium. The relationship between U and the intensity scintillation index is discussed.

In Section V, asymptotic results are derived for weak and strong scattering. The weak-scatter results were, of course, already deduced in Paper I. Under conditions of strong scattering, the asymptotic forms are critically dependent on the power-law spectral index, ν . [The three-dimensional SDF has the form $C_s q^{-(2\nu+1)}$ for $q \gg q_0$].*

When $\nu < 1.5$, simple asymptotic formulas can be derived for the autocorrelation function of intensity. Indeed, the S_4 scintillation index approaches unity from below and $\langle II' \rangle - 1 = \exp \{-2D(y)\}$, where $D(y)$ is the phase structure function. This behavior is demonstrated for a special case of one-dimensional scattering in which an exact result has been obtained.

When $\nu \geq 1.5$, however, the intensity statistics retain an explicit dependence on the Fresnel parameter Z , and S_4 can exceed unity. This particular finding is evidently new. In effect, the behavior of the scattering for the more steeply sloped spectra ($1.5 \leq \nu < 2.5$) is transitional between that for a power-law environment in which large-scale structures are strongly suppressed by Fresnel filtering ($0.5 < \nu < 1.5$) and a medium dominated by a single scale size where strong focusing can occur.

*The form of the three-dimensional index $q^{-\alpha}$ where $\alpha = 2\nu + 1$ is largely historical. The corresponding one-dimensional spectral index is $\alpha - 2 = 2\nu - 1$. For scintillation studies, it is the one-dimensional phase spectral index, p , that can be measured. In terms of ν , $p = 2\nu$. Thus, if $\nu = 1.5$, $\alpha = 4$ and $p = 3$.

In Section VI the results are summarized and compared to the earlier results (e.g., Salpeter, 1967) that apply strictly to scattering in a medium dominated by a single scale size, as opposed to a power-law continuum of scale sizes. A simple formula for the intensity coherence time is derived that fully accounts for angle effects in a highly anisotropic medium.

III GENERAL RESULTS FOR A GAUSSIAN PHASE SCREEN

Mercier (1962) developed a mathematically cumbersome but general formula for the second-order statistics of the intensity fluctuations of a wavefield as it propagates away from a phase-changing screen. A more compact formulation was developed by Gochelashvily and Shishov (1971). Alternative derivations of the Gochelashvily and Shishov result have been presented by Taylor (1972) and Rumsey (1975). Here we review only the principal results.

In the phase-screen model, a phase perturbation, $\delta\phi(\vec{\rho})$, is imparted to a wavefield at some plane, say $z = z_0$. Diffraction effects cause intensity fluctuations (scintillation) to develop as the wavefield propagates beyond $z = z_0$. The structure of intensity scintillation is characterized by the correlation function

$$R_I(\vec{\Delta\rho}; z) = \langle I(\vec{\rho}, z) I(\vec{\rho}', z) \rangle, \quad (1)$$

where $I(\vec{\rho}, z) = |u(\vec{\rho}, z)|^2$, $u(\vec{\rho}, z)$ is the complex wavefield at z , and the angle brackets denote mathematical expectation.

The free-space propagation of $R_I(\vec{\Delta\rho}; z)$ from z_0 to z can be calculated from the integral expression [the Gochelashvily and Shishov (1971) result]:

$$R_I(\vec{\Delta\rho}, z) = \frac{k}{z} \iiint M_4(\vec{\alpha}, \vec{\Delta\rho} - \vec{\beta}, 0; z_0) \exp \{-i\vec{\alpha} \cdot \vec{\beta} k/z\} d\vec{\alpha} d\vec{\beta} \quad (2)$$

where $k = 2\pi/\lambda$ and $\tilde{z} = z - z_0$. The quantity $M_4(\vec{\alpha}^{(2)}, \vec{\alpha}^{(3)}, \vec{\alpha}^{(4)}; z_0)$ is the fourth-order coherence function of the wavefield in the plane $z = z_0$, evaluated in terms of the variables

$$\vec{\alpha}^{(2)} = \frac{1}{2}[\vec{\rho}^{(1)} - \vec{\rho}^{(2)} - \vec{\rho}^{(3)} + \vec{\rho}^{(4)}] \quad (3a)$$

$$\vec{\alpha}^{(3)} = \frac{1}{2}[\vec{\rho}^{(1)} + \vec{\rho}^{(2)} - \vec{\rho}^{(3)} - \vec{\rho}^{(4)}] \quad (3b)$$

$$\vec{\alpha}^{(4)} = \frac{1}{2}[\vec{\rho}^{(1)} - \vec{\rho}^{(2)} + \vec{\rho}^{(3)} - \vec{\rho}^{(4)}] \quad (3c)$$

Two assumptions are used in deriving Eq. (2). First, the scattering of the principal components of the angular spectrum must be confined to a narrow cone about the direction of the reference wave. Under this assumption, the propagation effects are governed by the free-space form of the parabolic wave equation

$$\frac{\partial u(\vec{\rho}, z)}{\partial \ell} = -i \frac{1}{2k} \nabla_T^2 u(\vec{\rho}, z) \quad , \quad (4)$$

where $\partial u / \partial \ell$ is the directional derivative along the propagation path (Rino, 1978). Second, M_4 cannot depend on the "centroid" variable $\vec{\alpha}^{(1)} = \frac{1}{2}[\vec{\rho}^{(1)} + \vec{\rho}^{(2)} + \vec{\rho}^{(3)} + \vec{\rho}^{(4)}]$. That is, M_4 is statistically homogeneous, or independent of where it is measured. Note that this neither implies nor requires that lower order moments be statistically homogeneous.

To evaluate M_4 , the phase-screen model is used to obtain

$$M_4(\vec{\rho}^{(1)}, \vec{\rho}^{(2)}, \vec{\rho}^{(3)}, \vec{\rho}^{(4)}; z_0) = \langle \exp\{i[\delta\phi(\vec{\rho}_1) - \delta\phi(\vec{\rho}_2) + \delta\phi(\vec{\rho}_3) - \delta\phi(\vec{\rho}_4)]\} \rangle. \quad (5)$$

If $\delta\phi(\vec{\rho})$ is a zero-mean gaussian field, then Eq. (5) can be evaluated by using the well-known result

$$\langle \exp \{x\} \rangle = \exp \left\{ - \frac{\langle x^2 \rangle}{2} \right\} \quad . \quad (6)$$

The final form of Eq. (5) can then be conveniently written in terms of the phase structure function,

$$D(\vec{\Delta\rho}_{ij}) = \langle [\delta\phi(\rho^{(i)}) - \delta\phi(\rho^{(j)})]^2 \rangle \quad , \quad (7)$$

as

$$M_4(\vec{\rho}^{(1)}, \vec{\rho}^{(2)}, \vec{\rho}^{(3)}, \vec{\rho}^{(4)}; z_0) = \exp\left\{\frac{1}{2}[D(\Delta\vec{\rho}_{12}) - D(\Delta\vec{\rho}_{13}) + D(\Delta\vec{\rho}_{14}) - D(\Delta\vec{\rho}_{24}) + D(\Delta\vec{\rho}_{34}) + D(\Delta\vec{\rho}_{23})]\right\} \quad (8)$$

By making the appropriate substitutions from Eq. (3), it follows immediately that M_4 does not depend on $\vec{\alpha}^{(1)}$. Moreover,

$$M_4(\vec{\alpha}^{(2)}, \vec{\alpha}^{(3)}, 0; z_0) = \exp\left\{D(\vec{\alpha}^{(2)}) + D(\vec{\alpha}^{(3)}) - \frac{1}{2}D(\vec{\alpha}^{(2)} + \vec{\alpha}^{(3)}) - \frac{1}{2}D(\vec{\alpha}^{(2)} - \vec{\alpha}^{(3)})\right\} \quad (9)$$

Thus, the homogeneity of M_4 follows from the assumption that the structure function depends only on the difference variable $\vec{\Delta\rho}_{ij}$. It will be shown, however, that because of the symmetry of the variables in Eq. (9), Eq. (2) is well defined even if $D(\vec{\Delta\rho})$ does not exist.

The spectral domain version of Eq. (2) takes a particularly simple form. By direct computation

$$\begin{aligned} \tilde{\Phi}_I(\vec{k}) &= \iint R_I(\vec{\Delta\rho}) \exp\{i\vec{k} \cdot \vec{\Delta\rho}\} d\vec{\Delta\rho} \\ &= \iint M_4(-\vec{k}, \vec{k}; 0, z_0) \exp\{-i\vec{k} \cdot \vec{\xi}\} d\vec{\xi} \quad (10) \end{aligned}$$

By substituting Eq. (9) into Eq. (10), one obtains

$$\tilde{\Phi}_I(\vec{k}) = \iint \exp\{-g(\vec{\xi}, \vec{k})\} \exp\{-i\vec{k} \cdot \vec{\xi}\} d\vec{\xi} \quad (11)$$

where

$$g(\vec{\xi}, \vec{\eta}) = D(\vec{\xi}) + D(\vec{\eta}) - \frac{1}{2}D(\vec{\xi} + \vec{\eta}) - \frac{1}{2}D(\vec{\xi} - \vec{\eta}) \quad (12)$$

which is the principal result that will be used in later sections.

To investigate the general behavior of $g(\vec{\xi}, \vec{\eta})$, the spectral representation,

$$D(\vec{\xi}) = \iint [1 - \cos(\vec{q} \cdot \vec{\xi})] \Phi_{\delta\phi}(\vec{q}) d\vec{q}/(2\pi)^2, \quad (13)$$

can be used where $\Phi_{\delta\phi}(\vec{q})$ is the two-dimensional phase SDF. If this form is substituted into Eq. (12), after some straightforward manipulations, one obtains the equivalent representation

$$g(\vec{\xi}, \vec{\eta}) = 8 \iint \Phi_{\delta\phi}(\vec{q}) \sin^2(\vec{\eta} \cdot \vec{q}/2) \sin^2(\vec{\xi} \cdot \vec{q}/2) \frac{d\vec{q}}{(2\pi)^2}. \quad (14)$$

From the form of Eq. (14), it follows that if $\Phi_{\delta\phi}(\vec{q}) \propto q^{-\alpha}$, then $g(\vec{\xi}; \vec{\eta})$ is well defined as long as $2 < \alpha < 6$. From Eq. (13), however, it is clear that the structure function itself is only defined for $2 < \alpha < 4$.

To carry the development one step further, consider the special case in which $|g(\vec{\xi}, \vec{\eta})| \ll 1$ for all significant $\vec{\eta}$ values. The necessary conditions are stated in Section V. By using the approximation $\exp\{-g\} \cong 1 - g$, substituting from Eq. (14), and finally using the identity $\sin^2 x = (1 - \cos x)/2$, it follows from Eq. (11) that

$$\Phi_I(\vec{\kappa}) \cong \delta(\vec{\kappa}) + 4\Phi_{\delta\phi}(\vec{\kappa}) \sin^2[\kappa^2 \tilde{z}/(2k)], \quad (15)$$

which is the well known weak-scatter result.

To summarize, in the gaussian phase-screen model, the intensity SDF can be computed by evaluating a single integral, Eq. (11). Only the phase structure function need be specified for the computation. However, the diffraction effects, in conjunction with the size distribution of the irregularities, are completely characterized by the function $g(\vec{\xi}, \vec{\kappa} \tilde{z}/k)$ [Eq. (12)], which is well defined independently of an outer-scale cutoff for phase SDFs of the form $q^{-\alpha}$ with $2 < \alpha < 6$.

IV SCATTER IN STRICT POWER-LAW ENVIRONMENTS

In Section III, we showed that the second-order statistics of intensity are completely characterized by the function $g(\vec{\xi}, \vec{\eta})$, as defined by Eq. (12) in terms of the phase structure function. If the power-law model in Paper I is used,

$$\Phi_{\delta\phi}(\vec{q}) = \frac{abC_p}{[q_o^2 + q^2]^{\nu+1/2}}, \quad (16)$$

where

$$C_p = r_e^2 \lambda^2 L \sec \theta C_s \quad (17)$$

and

$$q^2 = A' \kappa_x^2 + B' \kappa_x \kappa_y + C' \kappa_y^2. \quad (18)$$

The coefficients A' , B' , and C' are defined in Paper I (Eqs. 26a, 26b, and 26c). The three-dimensional irregularity SDF has the same form as Eq. (16), except that abC_p is replaced by C_s . The parameters a and b are axial ratios along and transverse to the principal irregularity axis. A complete discussion of this model is given in Rino and Fremouw (1977).

The general form of the two-dimensional phase SDF corresponding to Eq. (16) is

$$R_{\delta\phi}(y) = \frac{GC_p}{2\pi\Gamma(\nu + 1/2)} \left| \frac{q_o y}{2} \right|^{\nu-1/2} K_{\nu-1/2}(q_o y)/q_o^{2\nu-1}, \quad (19)$$

where the geometric factor G is defined by Eq. (10) in Paper I. Hereafter, we shall consider only isotropic irregularities ($a = b = 1$). The general case is easily retrieved by reintroducing G and interpreting y appropriately. By using the small-argument formula,

$$K_{\nu-1/2}(x) \sim 1/2 \Gamma(\nu - 1/2) |x/2|^{-(\nu-1/2)},$$

it is readily shown that

$$R_{\delta\phi}(0) = \frac{C_p \Gamma(\nu - 0.5)}{4\pi \Gamma(\nu + 0.5) q_o^{2\nu-1}} \quad (20)$$

For completeness, a one-dimensional phase screen for which

$$\Phi_{\delta\phi}(q) = \frac{C'_p}{[q_o^2 + q^2]^{p/2}}, \quad (21)$$

$$R'_{\delta\phi}(y) = \frac{C'_p}{\sqrt{\pi} \Gamma(p/2)} \left| \frac{q_o y}{2} \right|^{\frac{p-1}{2}} K_{\frac{p-1}{2}}(q_o y) / q_o^{p-1}, \quad (22)$$

and

$$R'_{\delta\phi}(0) = \frac{C'_p \Gamma(\frac{p-1}{2})}{2\sqrt{\pi} \Gamma(p/2) q_o^{p-1}} \quad (23)$$

will also be considered. The one-dimensional model is appropriate for propagation across highly elongated irregularities. The relationship between p and ν is $p = 2\nu$.

Whenever the phase autocorrelation function (ACF) exists, the phase structure function can be written as

$$D(\vec{\Delta\rho}) = 2[R_{\delta\phi}(0) - R_{\delta\phi}(\vec{\Delta\rho})] \quad (24)$$

Thus, from Eqs. (19) and (20),

$$D(y) = \frac{C_p \Gamma(\nu - 0.5)}{2\pi \Gamma(\nu + 0.5)} \left[\frac{1 - 2 \left| \frac{q_o y}{2} \right|^{\nu-1/2} K_{\nu-1/2}(q_o y) / \Gamma(\nu - 0.5)}{q_o^{2\nu-1}} \right] \quad (25)$$

and

$$D'(y) = \frac{C'_p \Gamma\left(\frac{p-1}{2}\right)}{\sqrt{\pi} \Gamma(p/2)} \left[\frac{1 - 2\left|\frac{q_0 y}{2}\right|^{(p-1)/2} K_{(p-1)/2}(q_0 y) / \Gamma\left(\frac{p-1}{2}\right)}{q_0^{p-1}} \right] \quad (26)$$

As q_0 approaches zero, the correlation functions $R_{\delta\phi}(y)$ and $R'_{\delta\phi}(y)$ become singular. However, for a limited range of ν and p values, $D(y)$ and $D'(y)$ remain well defined.

The limiting forms of $D(y)$ and $D'(y)$ as q_0 approaches zero are not immediate, since $\lim_{q_0 \rightarrow 0} D(y) = 0/0$. Thus, one applies L'Hospital's rule, whereby the limit is obtained as the limit of the ratio of the separate derivatives of the numerator and the denominator. The results are

$$\begin{aligned} \lim_{q_0 \rightarrow 0} D(y) &= \lim_{q_0 \rightarrow 0} \frac{C_p}{2\pi\Gamma(\nu + 0.5)} \left[\frac{2\left|\frac{q_0 y}{2}\right|^{\nu-1/2} K_{\nu-3/2}(q_0 y) y}{(2\nu - 1)q_0^{2\nu-2}} \right] \\ &= \frac{C_p}{2\pi} \frac{2\Gamma(1.5 - \nu)}{\Gamma(\nu + 0.5)(2\nu - 1)} |y|^{2\nu-1} \end{aligned} \quad (27)$$

and

$$\begin{aligned} \lim_{q_0 \rightarrow 0} D'(y) &= \lim_{q_0 \rightarrow 0} \frac{C'_p}{\sqrt{\pi}\Gamma(p/2)} \left[\frac{2\left|\frac{q_0 y}{2}\right|^{(p-1)/2} K_{(p-1)/2}(q_0 y) y}{(p-1)q_0^{p-2}} \right] \\ &= \frac{C'_p}{\sqrt{\pi}} \frac{2\Gamma\left(\frac{3-p}{2}\right)}{\Gamma(p/2)(p-1)} |y|^{p-1} \end{aligned} \quad (28)$$

It follows from Eqs. (27) and (28) that the respective structure functions are well defined as q_0 approaches zero, as long as $0.5 < \nu < 1.5$ and $1 < p < 3$.

However, it was shown in Section II that $g(\vec{\xi}, \vec{\eta})$ is well defined beyond $\nu = 1.5$. Similarly, $g'(\vec{\xi}, \vec{\eta})$ is well defined beyond $p = 3$. Hence, the most general limiting form cannot be deduced by substituting

Eq. (27) or (28) into Eq. (12). Two methods of avoiding this dilemma have been discussed by Rumsey (1975). The most straightforward procedure is to apply the finite q_0 form of $D(y)$ or $[D'(y)]$ and use L'Hospital's rule to evaluate the limiting form of $g(\vec{\xi}, \vec{\eta})$. Thus,

$$g(\vec{\xi}, \vec{\eta}) = \frac{C_p}{2\pi} \frac{\Gamma(\nu - 0.5)}{\Gamma(\nu + 0.5)} \left\{ 1 - \frac{1}{\Gamma(\nu - 0.5)} \left[2|q_0 y_1|^{1/2} K_{\nu-1/2}(q_0 y_1) \right. \right. \\ + 2|q_0 y_2|^{1/2} K_{\nu-1/2}(q_0 y_2) \\ - |q_0 y_3|^{1/2} K_{\nu-1/2}(q_0 y_3) \\ \left. \left. - |q_0 y_4|^{1/2} K_{\nu-1/2}(q_0 y_4) \right] \right\} / q_0^{2\nu-1}, \quad (29)$$

where $y_1 = |\vec{\xi}|$, $y_2 = |\vec{\eta}|$, $y_3 = |\vec{\xi} - \vec{\eta}|$, and $y_4 = |\vec{\xi} + \vec{\eta}|$. A similar expression can be generated for $g'(\vec{\eta}, \vec{\xi})$. To evaluate Eq. (29) and the corresponding expression for $g'(\vec{\eta}, \vec{\xi})$, L'Hospital's rule must be applied twice. The results of this computation are summarized in Table 1.

The results in Table 1 for $g(\vec{\xi}, \vec{\eta})$ agree with Rumsey's (1975) Eqs. (18) and (19) for $0.5 < \nu \leq 1.5$, but not for $1.5 < \nu < 2.5$. The limiting forms of $g(\vec{\xi}, \vec{\eta})$ for $\nu < 1.5$ and $\nu > 1.5$ cannot be simply combined. A term has evidently been omitted in Rumsey's computation for $\nu > 1.5$ ($\alpha > 4$). In any case, $\nu = 1.5$ marks an important transition in the scattering behavior, which will be shown in Section IV.

To illustrate the significance of these results, let $\vec{\xi} = C(\nu)^{-1/(2\nu-1)} \vec{\xi}'$ in Eq. (11) where

Table 1
LIMITING FORMS OF $g(\xi, \eta)$ AND $g'(\xi, \eta)$

Two-Dimensional $g(\xi, \eta)$	
$C \frac{p}{2\pi} \frac{\Gamma(1.5 - \nu)}{\Gamma(0.5 + \nu)(2\nu - 1)2^{2\nu-1}} \left[2 \xi ^{2\nu-1} + 2 \eta ^{2\nu-1} - \xi - \eta ^{2\nu-1} - \xi + \eta ^{2\nu-1} \right]$	$0.5 < \nu < 1.5$
$C \frac{p}{2\pi} \left[- \xi ^2 \log \xi - \eta ^2 \log \eta + \frac{1}{2} \xi - \eta ^2 \log \xi - \eta + \frac{1}{2} \xi + \eta ^2 \log \xi + \eta \right]$	$\nu = 1.5$
$C \frac{p}{2\pi} \frac{\Gamma(2.5 - \nu)}{\Gamma(0.5 + \nu)(2\nu - 1)(2\nu - 2)2^{2\nu-2}} \left[-2 \xi ^{2\nu-1} - 2 \eta ^{2\nu-1} + \xi - \eta ^{2\nu-1} + \xi + \eta ^{2\nu-1} \right]$	$1.5 < \nu < 2.5$
One-Dimensional $g^{(1)}(\xi, \eta)$	
$C' \frac{p}{\sqrt{\pi}} \frac{\Gamma(\frac{3-p}{2})}{\Gamma(p/2)(p-1)2^{p-1}} \left[2 \xi ^{p-1} + 2 \eta ^{p-1} - \xi - \eta ^{p-1} - \xi + \eta ^{p-1} \right]$	$1 < p < 3$
$2C' \frac{p}{\sqrt{\pi}} \left[- \xi ^2 \log \xi - \eta ^2 \log \eta + \frac{1}{2} \xi - \eta ^2 \log \xi - \eta + \frac{1}{2} \xi + \eta ^2 \log \xi + \eta \right]$	$p = 3$
$C' \frac{p}{\sqrt{\pi}} \frac{\Gamma(5/2 - p/2)}{\Gamma(p/2)(p-1)(p-2)2^{p-2}} \left[-2 \xi ^{p-1} - 2 \eta ^{p-1} + \xi - \eta ^{p-1} + \xi + \eta ^{p-1} \right]$	$3 < p < 5$

$$C(\nu) = \begin{cases} \frac{C_p}{2\pi} \frac{\Gamma(1.5 - \nu)}{\Gamma(0.5 + \nu)(2\nu - 1)2^{2\nu-1}} & 0.5 < \nu < 1.5 \\ \frac{C_p}{2\pi} & \nu = 1.5 \\ \frac{C_p}{2\pi} \frac{\Gamma(2.5 - \nu)}{\Gamma(0.5 + \nu)(2\nu - 1)(2\nu - 2)2^{2\nu-2}} & 1.5 < \nu < 2.5 \end{cases} \quad (30)$$

Then, with an appropriate substitution for \vec{q} ,

$$C(\nu)^{1/(\nu-0.5)} \Phi_I(\vec{q}') C(\nu)^{1/(2\nu-1)} = \iint \exp \left\{ -h(\vec{\xi}', \vec{\eta}' / k C(\nu)^{1/(\nu-0.5)} \vec{q}') \right\} \cos(\vec{\xi}' \cdot \vec{q}') d\vec{\xi}' \quad (31)$$

where

$$h(\vec{\xi}, \vec{\eta}) = \begin{cases} \pm 2|\vec{\xi}|^{2\nu-1} \pm 2|\vec{\eta}|^{2\nu-1} \mp |\vec{\xi} - \vec{\eta}|^{2\nu-1} \mp |\vec{\xi} + \vec{\eta}|^{2\nu-1} & \nu \neq 1.5 \\ -|\vec{\xi}|^2 \log |\vec{\xi}| - |\vec{\eta}|^2 \log |\vec{\eta}| + 1/2 |\vec{\xi} - \vec{\eta}|^2 \log |\vec{\xi} - \vec{\eta}| \\ + 1/2 |\vec{\xi} + \vec{\eta}|^2 \log |\vec{\xi} + \vec{\eta}| & \nu = 1.5 \end{cases} \quad (32)$$

The form of Eq. (31) suggests the definition

$$U = (\tilde{z}/k)^{\nu-0.5} C(\nu) = C_p Z^{\nu-0.5} F(\nu) \quad (33)$$

where

$$Z = \lambda \tilde{z} / 4\pi \quad (34)$$

and

$$F(\nu) = 2^{\nu-0.5} C(\nu) / C_p \quad (35)$$

The reason for introducing the new definitions, Z and $F(v)$, will become clear shortly.

It now follows that the second-order statistics of intensity are completely characterized by the integral

$$I(\vec{q}; U) = \iint \exp \left\{ -h \left(\vec{\zeta}, U \left(\frac{1}{v-0.5} \right) \vec{q} \right) \right\} \cos (\vec{\zeta} \cdot \vec{q}) d\vec{\zeta} \quad (36)$$

Similarly, for the one-dimensional phase screen

$$I'(q'; U') = \int \exp \left\{ -h' \left(\zeta, U' \left(\frac{2}{p-1} \right) q \right) \right\} \cos (\zeta q) d\zeta \quad (37)$$

completely specifies the second-order statistics of intensity, where

$$\begin{aligned} U' &= (\tilde{z}/k)^{(p-1)/2} C'(p) \\ &= C' Z^{(p-1)/2} F'(p) \quad , \end{aligned} \quad (38)$$

$$C'(p) = \begin{cases} \frac{C'_p}{\sqrt{\pi}} \frac{\Gamma \left(\frac{3-p}{2} \right)}{\Gamma(p/2)(p-1)2^{p-1}} & 1 < p < 3 \\ \frac{2C'_p}{\sqrt{\pi}} & p = 3 \\ \frac{C'_p}{\sqrt{\pi}} \frac{\Gamma(5/2 - p/2)}{\Gamma(p/2)(p-1)(p-2)2^{p-2}} & 3 < p < 5 \end{cases} \quad , \quad (39)$$

and

$$F'(p) = 2^{(p-1)/2} C'(p)/C'_p \quad (40)$$

The definition of $h'(\zeta, \eta)$ is readily deduced from Table 1.

The single parameter U determines the general behavior of the second-order intensity statistics. In Section V asymptotic results will be derived for weak-scatter conditions ($U \ll 1$) and strong scatter conditions ($U \gg 1$). Before doing so, however, it is instructive to compare the U parameter to the intensity scintillation index S_4

$$[S_4^2 = R_I(0; z) - 1] \quad .$$

Paper I showed that for isotropic irregularities

$$S_4^2 = C_p Z^{\nu-0.5} \left[\frac{\Gamma\left(\frac{2.5-\nu}{2}\right)}{2\sqrt{\pi} (\nu-0.5) \Gamma\left(\frac{\nu+0.5}{2}\right)} \right] \quad . \quad (41)$$

Thus, from Eq. (33),

$$S_4^2 = U \left[\frac{\Gamma\left(\frac{2.5-\nu}{2}\right)}{2\sqrt{\pi} \Gamma\left(\frac{\nu+0.5}{2}\right) F(\nu)} \right] \quad . \quad (42)$$

Similarly, for the one-dimensional phase-screen,

$$S_4^2 = U' \left[\frac{2\Gamma\left(\frac{5-p}{4}\right)}{\sqrt{\pi} (p-1) \Gamma\left(\frac{p+1}{4}\right) F'(p)} \right] \quad . \quad (43)$$

For a fixed value of ν (or p), S_4^2 and U admit the same functional dependence on C_p and Z . However, the functional dependences of S_4^2 and U on ν are radically different.

From Eq. (41) it follows that S_4^2 is a continuous monotonically increasing function of ν . This happens because as the slope of the phase spectrum steepens with C_p fixed, the spectral intensity of wavenumbers corresponding to the Fresnel radius rapidly increases relative to the smaller-scale structures. The parameter U , by comparison, is sharply

discontinuous at $\nu = 1.5$ [U' is discontinuous at $p = 3$]. This is illustrated in Figure 1, where S_4^2/U is plotted as a function of ν .

In Section III it was shown that Fresnel filtering suppresses the influence of large-scale structures in intensity scintillation as long as $\nu < 2.5$. The behavioral transition at $\nu = 1.5$ can be interpreted as a boundary between regimes in which large-scale structures are strongly suppressed ($\nu < 1.5$) and those in which they are weakly suppressed ($\nu > 1.5$). The asymptotic behavior of $R_I(\vec{\Delta\rho}; z)$ for large U is very different in these two regimes, as we show in Section V.




FIGURE 1 RATIO OF S_4 TO U vs. ν SHOWING DISCONTINUITY AT $\nu = 1.5$

V ASYMPTOTIC RESULTS FOR LARGE AND SMALL VALUES OF U

Following the work of Gochelashvily and Shishov (1971), a number of authors have derived asymptotic formulas for $\hat{\Phi}_I(\vec{k}; z)$ and/or $R_I(\vec{\Delta p}; z)$ by using series expansions of $q(\vec{\xi}; \vec{\eta})$ (e.g., Taylor and Infosino, 1976; Buckley, 1971a, 1971b). Difficulties can arise, however, if $g(\vec{\xi}, \vec{\eta})$ is defined in terms of an implicit structure function. For example, moments do not exist unless an inner-scale cutoff is introduced. To do so, however, does not give physically meaningful results, since the basic integral [Eq. (11)] is well defined independent of either the inner-scale or the outer-scale cutoff.

Proper treatment of this problem demands that series approximations be applied to Eq. (32) directly, as Rumsey (1975) did for the special case of $\nu = 1$ ($\alpha = 3$). First note that $h(\vec{\xi}, 0) = 0$ irrespective of ν . Hence, for sufficiently small \vec{q} ,

$$\exp \left\{ -h \left(\vec{\xi}, U^{\frac{1}{\nu-0.5}} \vec{q} \right) \right\} \cong 1 - h \left(\vec{\xi}, U^{\frac{1}{\nu-0.5}} \vec{q} \right) \quad (44)$$

Substituting Eq. (44) into Eq. (36) gives the so-called low-frequency approximation

$$\begin{aligned} I_{LF}(\vec{q}; U) &= \delta(\vec{q}) - \iint h \left(\vec{\xi}, U^{\frac{1}{\nu-0.5}} \vec{q} \right) \cos(\vec{q} \cdot \vec{\xi}) d\vec{\xi} \\ &= \delta(\vec{q}) + 4 \left(C_p C(\nu)^{-1} \right) q^{-(2\nu+1)} \sin^2 \left(U^{\frac{1}{\nu-0.5}} q^2 / 2 \right) \quad (45) \end{aligned}$$

Note that the product $C_p C(\nu)^{-1}$ is independent of C_p .

The low-frequency approximation is valid for $q \ll \tilde{q} = U^{\frac{1}{\nu-0.5}}$. This is essentially equivalent to Gochelashvily and Shishov's result

when $\nu = 4/3$ ($\alpha = 11/3$). The situation is more complicated when $\vec{q} \gg \vec{\eta}$. For $\nu \neq 1.5$ and $\vec{\eta} \gg \vec{\xi}$ the starting point is

$$h(\vec{\xi}, \vec{\eta}) \cong \pm 2|\xi|^{2\nu-1} \mp 2 \frac{(\nu-0.5)\xi^2 + 2(\nu-0.5)(\nu-1.5)(\vec{\xi} \cdot \hat{a}_\eta)^2}{\eta^{3-2\nu}} \quad (46)$$

where $\hat{a}_\eta = \vec{\eta}/|\vec{\eta}|$. Ordering the terms in Eq. (46) and retaining only the lowest order terms in ξ/η gives the result

$$h(\vec{\xi}, \vec{\eta}) \cong \begin{cases} 2|\xi|^{2\nu-1} & 0.5 < \nu < 1.5 \\ (2\nu-1)[\xi^2 + (2\nu-3)(\vec{\xi} \cdot \hat{a}_\eta)^2] \eta^{2\nu-3} & 1.5 < \nu < 2.5 \end{cases} \quad (47)$$

For $\nu < 1.5$, the high-frequency approximation takes the particularly simple form

$$I_{HF}(\vec{q}; U) = \iint \exp[-2|\xi|^{2\nu-1}] \cos(\vec{q} \cdot \vec{\xi}) d\vec{\xi} \quad (48)$$

It follows from Eq. (48) that $I_{HF}(\vec{q}; U)$ converges rapidly to zero for $\vec{q} \gg 1$. Thus, if $U \ll 1$, the bulk of the intensity SDF is contained in the low-frequency approximation. For $U \gg 1$, $\vec{\eta}$ is very small, and the bulk of the intensity SDF (except the singularity at $\vec{q} = 0$) is contained in the high-frequency approximation. Thus,

$$I(\vec{q}; U) \cong \begin{cases} I_{LF}(\vec{q}; U) & U \ll 1 \\ \delta(\vec{q}) + I_{HF}(\vec{q}; U) & U \gg 1 \end{cases} \quad (49)$$

This behavior is nicely illustrated by the one-dimensional model with $p = 2$, wherein $I(\vec{q}; U)$ can be evaluated exactly. Indeed, for $p = 2$

$$h'(\xi, \eta) = \begin{cases} 2|\xi| & |\xi| \leq |\eta| \\ 2|\eta| & |\xi| > |\eta| \end{cases} \quad (50)$$

By substituting Eq. (50) into Eq. (34), it is readily shown that

$$\begin{aligned}
 I'(q; U') &= \delta(q) + 2 \int_0^{U'^2 |q|} \exp \{-2|\xi|\} \cos(q\xi) d\xi \\
 &= \delta(q) + \left[\frac{8 \sin^2(U'^2 q^2/2)}{4 + q^2} + \frac{2q \sin(qU'^2 |q|)}{4 + q^2} \right] \exp \{-2U'^2 |q|\} \\
 &\quad + \frac{4}{4 + q^2} \left[1 - \exp \{-2U'^2 |q|\} \right]. \quad (51)
 \end{aligned}$$

For $U' \ll 1$, and q small, the first \sin^2 term dominates Eq. (51) in agreement with the low-frequency approximation. For $U' \gg 1$, the third term dominates and there is no further dependence on U' . The second term in the square brackets is significant only in the intermediate q range.

By using the high-frequency approximation it is easily shown that for large U

$$R_I(\xi) \cong \exp \{-2|\xi|^{2\nu-1}\} + 1. \quad (52)$$

In terms of the original variables

$$\begin{aligned}
 R_I(y) - 1 &\cong \exp \{-2C(\nu)y^{2\nu-1}\} \\
 &= \exp \{-D(y)\} \quad (53)
 \end{aligned}$$

[see Eq. (27)]. It follows that $S_4^2 = R_I(0) - 1$ approaches unity as U becomes arbitrarily large. Moreover, upon recalling Bramely's (1954) result for a gaussian phase screen,

$$\langle uu'^* \rangle = \exp \left\{ -\frac{1}{2}D(y) \right\}, \quad (54)$$

it follows from Eq. (53) that $\langle II' \rangle - 1 = |\langle uu'^* \rangle|^2$, again as U becomes arbitrarily large.

If $u = x + iy$, where x and y are uncorrelated, equal-variance gaussian processes, then it is well known that $I = |u|^2$ is Rayleigh distributed and $\langle II' \rangle - 1 = |\langle uu^* \rangle|^2$. Moreover, the phase of U is uniformly distributed. Now, while Eq. (53) alone cannot guarantee uniformly distributed phase (modulo 2π) and Rayleigh statistics for intensity, experimental results (Fremouw et al., 1978; Rino, 1978) suggest that the Rayleigh model is a good approximation.

For $\nu > 1.5$, the high-frequency approximation [see Eq. (47)] is considerably more complicated. In particular, $I_{HF}(\vec{q}; U)$ retains an explicit dependence on U such that numeric computations must be used. Intuitively, however, it is clear that as ν increases from 1.5 to 2.5, the intensity statistics tend to behave more like the medium is being dominated by a single scale size, even though strictly speaking the outer scale is not important until ν exceeds 2.5.

When a single scale size is dominant, strong focusing can occur (Pisareva, 1958), which causes S_4 to exceed unity. The form of $I_{HF}(\vec{q}; U)$ for $\nu > 1.5$ does not exclude this from happening in a power-law environment, although the conditions that lead to $S_4 > 1$ have generated some controversy (Rumsey, 1975; Taylor and Infosino, 1976).

VI SUMMARY AND DISCUSSION

In this paper the power-law phase-screen model has been generalized to accommodate strong scattering. The intensity SDF $\Phi_I(\vec{q})$ was derived in terms of a single integral expression $I(\vec{q}; U)$:

$$\Phi_I(\vec{q}) = C(\nu)^{-1/(\nu-0.5)} I\left[\vec{q} C(\nu)^{-1/(2\nu-1)}, U\right], \quad (55)$$

where

$$U = C_p Z^{\nu-0.5} F(\nu) \quad (56)$$

has the same functional form as the weak-scatter formula for the scintillation index, S_4 . [Recall that $C_p = r_e^2 \lambda^2 (L \sec \theta) C_s$; where C_s is the strength-of-turbulence as defined by Eq. (7) in Paper I.] In general

$$S_4^2 = \iint I(\vec{q}, U) d\vec{q} (2\pi)^2 - 1, \quad (57)$$

whereas for $U \ll 1$,

$$S_4^2 \cong C_p Z^{\nu-0.5} \left[\frac{\Gamma\left(\frac{2.5-\nu}{2}\right)}{2/\pi (\nu-0.5) \Gamma\left(\frac{\nu+0.5}{2}\right)} \right] \mathfrak{F}, \quad (58)$$

where \mathfrak{F} is defined by Eq. (34) in Paper I.

It was shown in Section IV that as long as ν does not lie in the range $1.4 \leq \nu < 1.5$, then $S_4^2 > U$, where S_4^2 is evaluated using Eq. (58). Thus, the weak-scatter condition $U \ll 1$ need not require that $S_4 \ll 1$. This is consistent with the experience of most experimentalists, namely that the weak-scatter S_4 formula accurately reproduces their data, even for moderately large scintillation levels.

As ν approaches 1.5, $F(\nu)$ approaches infinity. The $\nu = 1.5$ point marks an important transition from a scintillation behavior regime in which the contribution of structures larger than the Fresnel radius is

strongly suppressed ($\nu < 1.5$) to one in which the contribution is only weakly suppressed ($\nu > 1.5$). When $\nu < 1.5$, the phase-structure function, for example, is well defined in the limit as the outer-scale wavenumber approaches zero.

In Section V an asymptotic analysis of $I(\vec{q}, U)$ was performed for large and small q values. For $U \ll 1$, the low-frequency approximation

$$\Phi_I(\vec{q}) \cong \delta(\vec{q}) + 4C_p q^{-(2\nu+1)} \sin^2(Zq^2) \quad (59)$$

accounts for all significant Fourier components. As U increases, the high-frequency approximation accounts for all the significant Fourier components, except the delta function at $\vec{q} = 0$. The form of the high-frequency approximation, however, is very critically dependent on ν .

When $\nu < 1.5$, the high-frequency approximation is independent of U . Thus, under conditions of strong scattering, the intensity statistics converge; ultimately there is no further change with increasing Z . When this happens, a simple analytic form for $\Phi_I(\vec{q})$ cannot be obtained. Nonetheless, the intensity correlation function takes the particularly simple form

$$\langle II' \rangle = \exp \{-D(y)\} \quad (60)$$

where

$$D(y) = G \frac{C_p}{2\pi} \left[\frac{2\Gamma(1.5-\nu)}{\Gamma(\nu+0.5)(2\nu-1)2^{2\nu-1}} \right] |y|^{2\nu-1} \quad (61)$$

To evaluate the temporal autocorrelation function of intensity, y is replaced by $v_{\text{eff}} \delta t$, where v_{eff} is defined by Eq. (13) in Paper I. Thus, if τ_I is defined to be the time separation to achieve an intensity decorrelation of e^{-1} , then

$$\tau_I = v_{\text{eff}} \left[G \frac{C_p}{2\pi} \left(\frac{2\Gamma(1.5-\nu)}{\Gamma(\nu+0.5)(2\nu-1)2^{2\nu-1}} \right) \right]^{\frac{1}{2\nu-1}} \quad (62)$$

The formula becomes meaningless as ν approaches 1.5. For $\nu \geq 1.5$, no simple formula such as Eq. (60), exists and numeric computations must be employed. Indeed, in that case τ_I will depend on Z , as it does under conditions of weak scatter, in which there is also no simple formula.

It is instructive to compare these results to the earlier results for a gaussian phase screen with a gaussian ACF of the form $\exp \{-\Delta \rho^2 / r_0^2\}$. For the gaussian ACF, the rms phase φ_0 can be computed from the expression

$$\varphi_0^2 = r_e^2 \lambda^2 (L \sec \theta) G \langle \Delta N_e^2 \rangle 2\pi^{-3/2} r_0 \quad . \quad (63)$$

Under conditions of weak, isotropic scattering ($\varphi_0 \ll 1$)

$$S_4^2 = 2\varphi_0^2 \left\{ 1 - \cos^2 \left(2\lambda \tilde{z} / (\pi r_0^2) \right) \right\} \quad . \quad (64)$$

If $\lambda z / r_0^2 \ll 1$,

$$S_4^2 \cong 2\varphi_0^2 \left(\frac{2\lambda \tilde{z}}{\pi r_0^2} \right)^2 \quad , \quad (65)$$

which is similar to Eq. (58). If, on the other hand, $\lambda \tilde{z} / r_0^2 \gg 1$,

$$S_4^2 \cong 2\varphi_0^2 \quad . \quad (66)$$

Moreover, Mercier (1962) has shown that the intensity statistics are Rician in this limit. By comparison, in a strict power-law environment, λz cannot be increased indefinitely without S_4 approaching unity. Put another way, in a strict power-law environment one always encounters strong scattering with increasing distance from the scattering region before S_4 saturates independent of \tilde{z} .

Under conditions of strong scattering, the relationship $\langle II' \rangle - 1 = \exp \{-2D(y)\}$ where

$$D(y) = \varphi_0^2 \left(1 - \exp \{-y^2 / r_0^2\} \right) \quad (67)$$

will ultimately apply. However, it is known that when $\varphi_0 > 1$ and

$$\varphi_0 \frac{\tilde{z}}{k} Q_0^2 \sim 1, \quad (68)$$

where Q_0^4 is the fourth-order moment of the gaussian spectrum, the scintillation index S_4 exceeds unity (Salpeter, 1967). This is the strong focusing effect. This phenomenon has a counterpart in a power-law medium. However, since neither φ_0 nor Q_0 exist, the strong focusing condition cannot be simply characterized as with Eq. (68).

The scintillation behavior in a gaussian ACF environment is conveniently summarized in diagram form, as was done initially by Cohen et al. (1967). Figure 2, the usual representation, is reproduced from Singleton (1970). Regions I and II correspond to the near and far zones, respectively. Region III is the "fully modulated" or saturation region where S_4 is unity and the relationship $\langle II' \rangle - 1 = \exp \{-2D(y)\}$ holds. Region IV is the strong focusing region where $S_4 > 1$.

In a strict power-law environment, by comparison, there is no characteristic scale size akin to r_0 , and the magnitude of the rms phase is irrelevant to the scattering behavior. Contours of constant S_4 have the form $U = \text{const.}$, or, from Eq. (56),

$$C_p z^{\nu-0.5} = \text{const.} \quad (69)$$

Thus, if one were to make a diagram similar to the Cohen-Singleton diagram for a power-law medium, the phase strength-of-turbulence, C_p , would replace φ_0 and the Fresnel area $\lambda z/4\pi$ would replace $\lambda z/r_0^2$.

The degree to which Region II is present depends critically on the spectral index ν . If $\nu < 1.5$, S_4 does not significantly exceed unity. For $\nu \geq 1.5$, however, S_4 values substantially larger than 1 do occur. When this happens, the contours are still given by Eq. (69). Thus, the scattering behavior in a strict power-law environment can be similar to that predicted by the Cohen-Singleton diagram for $\lambda z/r_0^2 \ll 1$.

In fact, if a finite outer-scale is introduced, then the Cohen-Singleton diagram for a power-law medium is similar to that of a gaussian




FIGURE 2 CURVES OF CONSTANT S_4 ON A
PLOT OF $\lambda z/r_0^2$ vs. ϕ_0 --TAKEN FROM
SINGLETON (1970)

medium with the outer scale playing the role of r_0 . Because the outer scale is at best indeterminant in the ionosphere, however, such an extrapolation is not useful for data interpretation.

REFERENCES

- Bramely, E. N., "The Diffraction of Waves by an Irregular Refracting Medium," Proc. Roy. Soc. A., Vol. 225, p. 515, 1954.
- Buckley, R., "Diffraction by a Random Phase Screen with Very Large R.M.S. Phase Deviation I. One-Dimensional Screen," Aust. J. Phys., Vol. 24, p. 351, 1971a.
- Buckley, R., "Diffraction by a Random Phase Screen with Very Large R.M.S. Phase Deviation II. Two-Dimensional Screen," Aust. J. Phys., Vol. 24, p. 373, 1971b.
- Cohen, M. H., E. J. Gunermann, H. E. Hardebeck, and L. E. Sharp, "Interplanetary Scintillations II. Observations," Astrophys. J., Vol. 147, p. 449, 1967.
- Fante, R. L., "Electromagnetic Beam Propagation in Turbulent Media," Proc. IEEE, Vol. 63, No. 12, p. 1669, 1975.
- Fremouw, E. J., R. L. Leadabrand, R. C. Livingston, M. D. Cousins, C. L. Rino, B. C. Fair, and R. A. Long, "Early Results from the DNA Wideband Satellite Experiment--Complex-Signal Scintillation," Rad. Sci., Vol. 13, No. 1, p. 167, 1978.
- Gochelashvily, K. S. and V. I. Shishov, "Multiple Scattering of Light in a Turbulent Medium," Optica Acta, Vol. 18, No. 10, p. 767, 1971.
- Mercier, R. P., "Diffraction by a Screen Causing Large Random Phase Fluctuations," Proc. Cambr. Phil. Soc., Vol. 58, p. 382, 1962.
- Pisareva, V. V., "The Diffraction of Radio Waves by Random Nonuniformities, and Intensity Variations of Solar and Cosmic Radio Emission," Astronomy J. (USSR), Vol. 2, p. 97, 1958.
- Rino, C. L., "Iterative Methods for Treating the Multiple Scattering of Radio Waves," J. Atmos. Terres. Phys., Vol. 40, p. 1011, 1978.
- Rino, C. L. and E. J. Fremouw, "The Angle Dependence of Singly Scattered Wavefields," J. Atmos. Terr. Phys., Vol. 39, p. 859, 1977.
- Rino, C. L. and S. J. Matthews, "On the Interpretation of Ionospheric Scintillation Data Using a Power-Law Phase Screen Model--Weak Scatter," DNA4606T, Topical Report 2, Contract DNA001-77-C-0220, SRI Project 6434, SRI International, Menlo Park, California, 1978.

- Rumsey, V. H., "Scintillations Due to a Concentrated Layer with a Power-Law Turbulence Spectrum," Radio Sci., Vol. 10, No. 1, p. 107, 1975.
- Salpeter, E. E., "Interplanetary Scintillations I. Theory," Astrophys. J., Vol. 147, p. 433, 1967.
- Singleton, D. G., "Saturation and Focusing Effects in Radio-Star and Satellite Scintillations," J. Atmos. Terr. Phys., Vol. 32, p. 187, 1970.
- Taylor, L. S., "Scintillation of Randomized Electromagnetic Fields," J. of Mathematical Phys., Vol. 13, p. 590, 1972.
- Taylor, L. S. and C. J. Infosino, "On the Strong Phase Screen Theory of Ionospheric Scintillations," Radio Sci., Vol. 11, No. 5, p. 459, 1976.

DISTRIBUTION LIST

DEPARTMENT OF DEFENSE

Assistant to the Secretary of Defense
ATTN: Executive Assistant

Command & Control Technical Center
Department of Defense
ATTN: C-650

Defense Advanced Rsch. Proj. Agency
ATTN: T10

Defense Communications Agency
ATTN: Code 480
ATTN: Code 810, J. Barna
ATTN: Code 101B
ATTN: Code R1033, M. Raffensperger

Defense Communications Engineer Center
ATTN: Code R820
ATTN: Code R410, J. McClean

Defense Documentation Center
12 cy ATTN: DD

Defense Nuclear Agency
ATTN: DDST
ATTN: RAAE
ATTN: STVL
4 cy ATTN: TITL

Field Command
Defense Nuclear Agency
ATTN: FCPR

Field Command
Defense Nuclear Agency
ATTN: FCPRL

Interservice Nuclear Weapons School
ATTN: TTV

Joint Chiefs of Staff
ATTN: J-3, WWMCCS Evaluation Office

Joint Strat. Tgt. Planning Staff
ATTN: JLTW-2

National Security Agency
ATTN: R52, J. Skillman

NATO School (SHAPE)
ATTN: U.S. Documents Officer

Under Secy. of Def. for Rsch. & Engrg.
ATTN: Strategic & Space Systems (OS)

WWMCCS System Engineering Org.
ATTN: R. Crawford

DEPARTMENT OF THE ARMY

BMD Advanced Technology Center
Department of the Army
ATTN: ATC-T, M. Capps

DEPARTMENT OF THE ARMY (Continued)

Harry Diamond Laboratories
Department of the Army
ATTN: DELHD-N-P, F. Wimenitz
ATTN: DELHD-I-TL, M. Weiner
2 cy ATTN: DELHD-N-P

U.S. Army Comm-Elec. Engrg. Instal. Agency
ATTN: CC-EMEO-PED, G. Lane

U.S. Army Foreign Science & Tech. Ctr.
ATTN: DRXST-SD

U.S. Army Satellite Comm. Agency
ATTN: Document Control

U.S. Army TRADOC Systems Analysis Activity
ATTN: ATAA-PL

DEPARTMENT OF THE NAVY

Naval Electronic Systems Command
ATTN: Code 501A
ATTN: PME 106-4, S. Kearney
ATTN: PME 106-13, T. Griffin

Naval Intelligence Support Ctr.
ATTN: NISC-50

Naval Ocean Systems Center
ATTN: Code 532

Naval Research Laboratory
ATTN: Code 7500, B. Wald
ATTN: Code 6700, T. Coffey
ATTN: Code 7580
ATTN: Code 7175, J. Johnson

Naval Surface Weapons Center
ATTN: Code F31

Strategic Systems Project Officer
Department of the Navy
ATTN: NSP-2141
ATTN: NSSP-2722, F. Wimberly

DEPARTMENT OF THE AIR FORCE

Air Force Avionics Laboratory
ATTN: AAD, A. Johnson

Air Force Geophysics Laboratory
ATTN: OPR-1, J. Ulwick
ATTN: PHP, J. Mullen
ATTN: PHP, J. Aarons
ATTN: PHI, J. Buchau

Air Force Technical Applications Center
ATTN: TN

Air Force Weapons Laboratory
ATTN: DYC, J. Barry
ATTN: SUL
ATTN: DYC, J. Kamm
ATTN: DYT, M. Fry

DEPARTMENT OF THE AIR FORCE (Continued)

Foreign Technology Division, AFSC
ATTN: NIIS Library

Rome Air Development Center, AFSC
ATTN: TSLD

Space & Missile Systems Organization
Air Force Systems Command
ATTN: MNNL, S. Kennedy

Space & Missile Systems Organization
Air Force Systems Command
ATTN: SKA, M. Clavin

Strategic Air Command
Department of the Air Force
ATTN: NRT
ATTN: XPFS, B. Stephan
ATTN: ADWATE, B. Bauer

OTHER GOVERNMENT AGENCIES

Central Intelligence Agency
ATTN: OSI/PSTD, Rm. 5 F 19

Institute for Telecommunications Sciences
National Telecommunications & Info. Admin.
ATTN: W. Utlaut

DEPARTMENT OF ENERGY CONTRACTORS

Lawrence Livermore Laboratory
University of California
ATTN: Technical Information Department

Los Alamos Scientific Laboratory
ATTN: R. Jeffries
ATTN: J. Wolcott
ATTN: J. Zinn

Sandia Laboratories
ATTN: W. Brown
ATTN: D. Dahlgren
ATTN: T. Wright

DEPARTMENT OF DEFENSE CONTRACTORS

Aerospace Corp.
ATTN: N. Stockwell
ATTN: J. Carter
ATTN: I. Garfunkel
ATTN: R. Slaughter

Berkely Research Associates, Inc.
ATTN: J. Workman

Boeing Co.
ATTN: D. Murray
ATTN: G. Keister

Charles Stark Draper Lab, Inc.
ATTN: J. Gilmore
ATTN: D. Cox

Computer Sciences Corp
ATTN: J. Spoor
ATTN: C. Nail

DEPARTMENT OF DEFENSE CONTRACTORS (Continued)

Cornell University
Department of Electrical Engineering
ATTN: D. Farley, Jr.

Electrospace Systems, Inc.
ATTN: P. Phillips

ESL, Inc.
ATTN: C. Prettie
ATTN: J. Marshall

General Electric Co.
ATTN: F. Reibert

General Electric Company-TEMPO
Center for Advanced Studies
ATTN: W. Knapp
ATTN: DASIAC

General Research Corp.
ATTN: J. Garbarino
ATTN: J. Ise, Jr.

Geophysical Institute
ATTN: Technical Library

University of Illinois
ATTN: K. Yeh

Institute for Defense Analyses
ATTN: E. Bauer

International Tel. & Telegraph Corp.
ATTN: Technical Library

Jaycor
ATTN: S. Goldman

Johns Hopkins University
Applied Physics Lab
ATTN: T. Potemra
ATTN: Document Librarian

Kaman Sciences Corp.
ATTN: T. Meagher

Linkabit Corp.
ATTN: I. Jacobs

University of Lowell Rsch. Foundation
ATTN: K. Bibl

M.I.T. Lincoln Lab
ATTN: D. Towle

McDonnell Douglas Corp.
ATTN: Technical Library Services

Mission Research Corp.
ATTN: R. Bogusch
ATTN: R. Hendrick
ATTN: R. Kilb
ATTN: D. Sappenfield
ATTN: E. Fajen

DEPARTMENT OF DEFENSE CONTRACTORS (Continued)

Mitre Corp.

ATTN: J. Wheeler
ATTN: W. Foster
ATTN: W. Hall

R & D Associates

ATTN: W. Karzas
ATTN: R. Lelevier
ATTN: B. Gabbard
ATTN: C. MacDonald

Rand Corp.

ATTN: C. Crain
ATTN: E. Bedrozian

DEPARTMENT OF DEFENSE CONTRACTORS (Continued)

Science Applications, Inc.

ATTN: L. Linson
ATTN: D. Sachs
ATTN: D. Hamlin

SRI International

ATTN: D. McDaniels
ATTN: R. Hake, Jr.
ATTN: R. Leadabrand
ATTN: W. Chestnut
ATTN: V. Gonzales
ATTN: C. Rino

Technology International Corp.

ATTN: W. Boquist